

Clinical Paper
Reconstructive Surgery

Temporomandibular joint reconstruction with medial femoral condyle osseocartilaginous flap: a case series

L. Xia^{1,2,3,4,a}, B. Jie^{1,2,3,4,a},
Y. Zhang^{1,2,3,4}, J. An^{1,2,3,4},
L. Zheng^{1,2,3,4}, Y. He^{1,2,3,4}

¹Department of Oral and Maxillofacial Surgery, Peking University School and Hospital of Stomatology, Beijing, PR China; ²National Clinical Research Center for Oral Diseases, Beijing, PR China; ³National Engineering Laboratory for Digital and Material Technology of Stomatology, Beijing, PR China; ⁴Beijing Key Laboratory of Digital Stomatology, Beijing, PR China

L. Xia, B. Jie, Y. Zhang, J. An, L. Zheng, Y. He: Temporomandibular joint reconstruction with medial femoral condyle osseocartilaginous flap: a case series. *Int. J. Oral Maxillofac. Surg.* 2021; 50: 604–609. © 2020 International Association of Oral and Maxillofacial Surgeons. Published by Elsevier Ltd. All rights reserved.

Abstract. Reconstruction for large-scale temporomandibular joint (TMJ) defects can be challenging. We have used the medial femoral condyle (MFC) osseocartilaginous flap for repair of TMJ defects. The aim of this paper was to describe our technique and to present the preliminary results. The MFC osseocartilaginous flap was used as a free vascularized graft for TMJ defect in four patients who had undergone resection for benign tumor of the TMJ region ($n = 2$) or TMJ ankylosis ($n = 2$). A computer-assisted technique was used in all cases. Symmetry of the mandible was objectively evaluated by postoperative computed tomography. Complications were recorded during follow-up visits. Lower extremity functional status was assessed by the Lower Extremity Functional Scale (LEFC) questionnaire. The MFC osseocartilaginous flap success rate was 100%; bony union was obtained in all four patients, and normal occlusion was achieved within 6–11 months after surgery. No flap-related complications occurred. All patients were satisfied with their postoperative facial symmetry and oral function. The LEFC score ranged from 72 to 80, indicating normal lower extremity function. Vascularized MFC osseocartilaginous flap appears to be a reliable option for reconstruction of large-scale TMJ defects.

Key words: temporomandibular joint; temporomandibular ankylosis; mandibular reconstruction; microsurgical free flaps; femur.

Accepted for publication 21 September 2020
Available online 9 October 2020

The temporomandibular joint (TMJ) plays an important role in mastication, speech and facial aesthetics, hence accurate reconstruction for large-scale defects of the TMJ is essential¹. Traditional surgical

approaches include non-vascularized autogenous reconstruction, vascularized autogenous reconstruction, and alloplastic reconstruction. Each method has limitations; for example, bone grafting is often

associated with severe resorption and vascularized bone flap surgery with a number

^a These authors contributed equally to this study.

of donor-site morbidities, while alloplastic prosthesis is either not available or not affordable in many countries. Hence, it is important to identify alternative methods.

The medial femoral condyle (MFC) flap, which incorporates cortical bone, cancellous bone, periosteum, muscle, tendon, cartilage, and skin, is well suited for complex reconstructions². In addition, donor-site morbidity is reported to be acceptable, with a good level of postoperative function^{3,4}. However, the vascularized MFC flap, carrying with it articular cartilage tissue, has not been previously used for reconstruction for large-scale TMJ defects. We have recently developed a technique for condyle reconstruction using the MFC osseocartilaginous flap and have used it to treat four patients at our hospital. The aim of this paper was to describe the technique and present the preliminary results.

Materials and methods

Between 2019 and 2020, we used the MFC osseocartilaginous flap to perform TMJ reconstruction for four patients. The patients (three females, one male) were in the age range of 40–59 years (mean, 49.8 years). The primary diagnoses included condylar osteochondroma (*n* = 1), condylar osteoma (*n* = 1), and type IV TMJ ankylosis (*n* = 2)⁵. The patients were followed up for periods ranging from 6 to 11 months (mean, 8.5 months). **Table 1** lists the demographic and clinical characteristics of the four patients. Written informed consent was obtained from patients before their surgeries.

Virtual surgical planning

All patients underwent spiral computed tomographic (CT) scanning of the maxillofacial and knee joint regions (field of view, 20 cm; pitch, 1.0; slice, 0.75 mm; 120Y280 mA). The scan data were imported into ProPlan CMF software (Materialise, Leuven, Belgium), to produce a direct three-dimensional (3D) model. The resection margin was determined according to the clinical and 3D radiographic findings, and the condylectomy was simulated using software.

An MFC osseocartilaginous flap was planned based on the defect shape and size, and arranged to reconstruct the TMJ region on the 3D radiographic image. To shape the bony flap to resemble the native condyle as much as possible, 3D patient-specific surgical templates were designed, based on mirror images of the unaffected side, and then fabricated using

computer-assisted design and manufacturing techniques (**Figs 1 and 2**).

Surgical procedure

A preauricular incision was used to expose the TMJ region, and osteoarthrectomy or condylectomy was performed from the top of the glenoid fossa to the residual mandibular ramus. Resection of the coronoid process was performed, if needed, to ensure that >40 mm of passive maximal mouth opening (MMO) was achieved. After condylectomy, the main surgical team (the authors) focused on the ipsilateral neck to identify and prepare adequate recipient vessels for microvascular anastomosis. Meanwhile, a second team harvested the ipsilateral osteochondral MFC flap, along with a pedicle containing the descending genicular artery. Because the transverse branch of the artery courses anteriorly towards the medial condyle and is invested intimately with the maximum range of the cartilage, it was preserved such that this convex cartilaginous surface could be harvested as a free vascularized flap. A pre-made surgical guide was used when necessary to confirm the shape of the flap before the osteotomies were performed on the femoral condyle (**Fig. 3**).

The flap was transferred to the head, where the pre-made guide was used to adjust the position of the flap in relation to the glenoid fossa and to adapt it to the native mandible before it was fixed in place with two miniplates. The temporal muscle myofascial flap was used to obliterate the soft tissue defect and also to separate the bony flap from the skull base. Microvascular anastomosis was completed under the operating microscope, and the wound was closed after placement of a suction drain (**Fig. 4**).

All the surgeries were performed by the same team, comprising experienced senior surgeons in the fields of reconstructive surgery.

Postoperative management

Patients were required to perform mouth opening exercises regularly over 3 months, starting at 1 week after the surgery. Facial symmetry was self-evaluated and scored by the patient on a scale of 0 to 10; the results were classified as good (8–10), fair (4–7), or poor (0–3).

Lower extremity functional status was assessed by postoperative knee radiographs and a validated condition-specific measure – the Lower Extremity Functional Scale (LEFC) questionnaire⁶. The items

Table 1. Demographic and clinical characteristics of the four patients, and the postoperative outcomes.

Patient	Sex	Age (years)	Affected side	Primary diagnosis	Size of defect (mm)	Size of bone flap (mm)	MMO (mm)	Self-evaluation of appearance (0–10)			Follow-up (months)
								LEFC score at T1 (0–80)	LEFC score at T2 (0–80)	LEFC score at T2 (0–80)	
1	F	46	Left	Condylar osteochondroma	40.7 × 22.5	50.3 × 32.1	30	9	15	80	11
2	M	40	Left	Condylar osteoma	34.6 × 16.8	51.5 × 31.9	36	7	13	72	9
3	F	54	Left	TMJ ankylosis	35.1 × 21.9	46.3 × 39.5	24	9	12	80	8
4	F	59	Right	TMJ ankylosis	33.6 × 22.8	45.5 × 36.9	36	8	17	73	6

F, female; LEFC, Lower Extremity Functional Scale; M, male; MMO, maximal mouth opening; TMJ, temporomandibular joint; T1, 1 week after surgery; T2, follow-up time.




Fig. 1. Virtual surgical planning. (A) Simulation of the bony flap on the three-dimensional radiographic image of the femur. (B) Arrangement of the bony flap to reconstruct the defect, using mirror images of the unaffected side for guidance. (C) Fabrication of individualized surgical templates of the bony flap.




Fig. 2. Transfer of the planned medial femoral condyle osseocartilaginous flap to the temporomandibular joint region. DGA, descending genicular artery.




Fig. 3. Harvesting of the medial femoral condyle osseocartilaginous flap. (A) Exposure of the medial femoral condyle with its articular cartilage surface. (B) Shaping of the bony flap according to the surgical template.




Fig. 4. The medial femoral condyle osseocartilaginous flap is transplanted to the defect. (A) The bony flap is positioned at the buccal side of the mandibular ramus, with the articular cartilage surface located in the glenoid fossa. (B) Bony suture with two miniplates, and microvascular anastomosis to the appropriate vessels in the recipient area.

in the questionnaire investigate the degree of difficulty in performing different physical activities. The LEFC score can range from 0 (maximal disability) to 80 (no disability).

Results

Postoperative hospital stay was 8–10 days. There were no intraoperative or

postoperative complications such as infection, haemorrhage, or facial nerve injury. MMO at the last follow-up ranged from 24 to 36 mm (mean, 31.5 mm). All patients had slight deviation of the mandible to the affected side during mouth opening. The facial symmetry achieved was self-evaluated as ‘good’ by three patients and as ‘fair’ by one patient.

Postoperative CT scans and 3D reconstructed images showed excellent anatomic restoration with the bony flaps. The flap success rate was 100% (4/4) for patency and achieving osseous union. There were no flap-related complications such as postoperative haematoma, vascular thrombosis, or flap necrosis. All patients achieved normal load-bearing occlusion within 6–11 months after surgery.




Fig. 5. A representative patient. (A) A 46-year-old woman was referred to us with complaint of swelling and discomfort in the left preauricular area. (B) The condyle was severely damaged by a tumor. (C) A condylectomy of >4 cm was performed. (D) At follow-up 9 months after surgery, the mandibular contour is intact. (E) There is a well-integrated bone graft without significant resorption. (F) Mouth opening is also satisfactory.

While all four patients experienced pain and numbness of skin on the anteromedial aspect of the knee region after surgery, the symptoms resolved without treatment within 3 months. LEFS score at the last follow-up ranged from 72 to 80 (mean, 76.3); two of the four patients scored 80, indicating normal level of function.

Fig. 5 presents a representative case.

Discussion

Traditional methods for reconstruction of large-scale TMJ defects include non-vascularized autogenous reconstruction, vascularized autogenous reconstruction and alloplastic reconstruction. Each method has disadvantages.

Alloplastic prosthesis, especially the custom-made total TMJ prosthesis, is a safe and predictable way to restore acquired TMJ defects^{7,8}; unfortunately, in several regions of the world such prosthesis is either not available or not affordable.

The greatest drawback of non-vascularized autogenous bone graft in TMJ reconstruction is the severe degree of bone resorption, especially in cases where a large bone defect has been reconstructed. Huang et al.⁹ found a high frequency of bone resorption with both the free coronoid process graft and the costochondral graft in adult patients undergoing condylar reconstruction. Although costochondral graft is considered suitable for children because of its intrinsic potential for growth, it is limited by the unpredictability of the growth pattern; resorption, failure to grow, and overgrowth have all been reported¹⁰.

In addition, our previous research¹¹ found that the heights of condyles reconstructed by transport distraction osteogenesis of the mandibular ramus were not stable in the long term, and the mean resorption rate was as high as 59.4%. Chen et al.¹² tackled the problem of shortage of blood supply using the superior half of the sternoclavicular joint, pedicled with the clavicular head of the sternocleidomastoid muscle and sternocleidomastoid branch of the superior thyroid artery; however, this approach was associated with high risk of donor-site morbidity, and the remaining clavicle had to be fixed with a thick titanium plate to avoid fracture.

These problems can be overcome with the use of vascularized bone flaps, which have lower bone resorption rates and superior bony quality¹³. The free fibula flap – a workhorse flap for mandibular reconstruction – is the first choice when bony length is required, as in reconstruc-

tion of subtotal or total mandibulectomy¹⁴. However, the fibula flap is more suitable for the reconstruction of a total mandibular defect and for defects that cross the midline, than for large-scale TMJ defects. It is very difficult to confirm whether the top of the transferred fibula is inserted precisely into the glenoid fossa when fossa exposure is inadequate. With the conventional free fibula flap there is no real cartilage-covered condyle. If the fibula head (with a branch of the anterior tibial artery) and the syndesmosis is taken for condyle reconstruction, a different situation results. The problems then include possible peroneal nerve damage and donor site morbidities such as ankle instability/stiffness, motor weakness or clawing of the great toe, and tendon exposure secondary to skin graft loss¹⁵.

The MFC flap has many advantages. It has been used for reconstruction at various head and neck sites, from the orbit, maxilla, and mandible, to the laryngeal and tracheal scaffolds^{16–18}. Lee et al.¹⁹ have used the MFC bone flap for treatment of malunion after subcondylar fracture. In addition, Zeman-Kuhnert et al.⁴ reported minimal donor site morbidity after harvesting MFC for maxillofacial indications; no patient in their study developed knee joint instability or limitation of range of motion.

The importance of subchondral blood supply for the nutrition of the chondrocytes in articulating cartilage is well known. Higgins et al.²⁰ showed that articular reconstruction performed using vascularized osteochondral flaps yields superior cartilage organization and architecture than reconstruction performed using non-vascularized osteochondral grafts. In addition, Benlidayi et al.²¹ showed in an experimental study that the bone resorption rate is greater in non-vascularized femur grafts than in vascularized osteoperiosteal femur flaps.

The usefulness of the vascularized MFC, harvested with the convex cartilaginous surface, was first demonstrated in 2008 in a patient requiring scaphoid reconstruction²². Subsequent studies have described the use of the MFC carrying cartilaginous tissue for carpal reconstruction²³, advanced Kienbock disease²⁴, and scaphoid and lunate reconstruction²⁵. Cadaver studies^{26,27} have shown that the descending genicular artery could provide good and reliable periosteal perfusion, suggesting that corticocancellous MFC might provide the benefits of vascularized bone for large intercalary nonunion defects that have conventionally been treated with fibula flaps.

Surgical repair of large TMJ defects is difficult mainly because of the complex tissue reconstructions required, including that of the condyle, ramus, and soft tissue. All of our patients had undergone large resections for tumor or extensive bony ankylosis. Free bone grafts will not be enough to repair such defects. Our technique of using the MFC osseocartilaginous flap for reconstruction of large-scale TMJ defects offers the following advantages: (1) the MFC flap contains sufficient cortical and cancellous bone to pack such a bony defect; (2) the articular cartilage tissue in the flap resembles the TMJ cartilage anatomically; and (3) preservation of the transverse branch of descending genicular artery prevents postoperative bone resorption of the grafts and ensures the success of functional rehabilitation.

In 2018, Enzinger et al.²⁸ performed mandibular condyle reconstruction using an osteochondral lateral femoral condyle flap, which is harvested from almost the same region as the flap we used. Our method had some notable advantages: (1) we combined preauricular and submandibular approaches to ensure better exposure, more extended resection, and more choices for internal fixation; (2) we chose the facial arterial as the recipient vessel to make microvascular anastomosis, and thus there is no need to fold the DGA for anastomosis with the temporal superficial artery; and (3) we aim to obtain a triangle-shaped bony flap, with the narrow side as condyle and the wider side as mandibular body, which allows reconstruction of relatively large defects.

To our knowledge, this is the first report of the use of the MFC flap, carrying articular cartilage, for TMJ reconstruction. The preliminary results suggest satisfactory restoration of mandibular height and facial symmetry, and good recovery of function (mastication, articulation, speech). However, the technique has limitations. The transplanted bone flap will inevitably thicken the affected side of the face, as was observed in one of our patients. Ideally, we should have investigated the histological transformation of osseocartilaginous flap; however, this was not done because of ethical considerations.

Vascularized MFC osseocartilaginous flap appears to be a reliable option for reconstruction of large-scale TMJ defects. Large prospective studies with long follow-up are needed to determine long-term outcomes and the risk of late lower extremity functional problems.

Funding

The article was supported by National Key R&D Program of China (2017YFB1104103), National Key R&D Program of China (2019YFF0302401) and the Program for New Clinical Techniques and Therapies of Peking University School and Hospital of Stomatology (grant PKUSSNCT-19A01). The funding agencies had no role in the study design, clinical procedures, analysis of the results, or preparation of the manuscript.

Competing interests

None.

Ethical approval

This study was approved by the ethics committee of Peking University School and Hospital of Stomatology (No. PKUS-SIRB-201949133).

Patient consent

All patients or their legal guardians have signed informed consents for publication of clinical photographs.

References

- Vega LG, Gonzalez-Garcia R, Louis PJ. Reconstruction of acquired temporomandibular joint defects. *Oral Maxillofac Surg Clin North Am* 2013;**25**:251–69.
- Banaszewski J, Gaggl A, Andruszko A. Medial femoral condyle free flap for head and neck reconstruction. *Curr Opin Otolaryngol Head Neck Surg* 2019;**27**:130–5.
- Mehio G, Morsy M, Cayci C, et al. Donor-Site Morbidity and Functional Status following Medial Femoral Condyle Flap Harvest. *Plast Reconstr Surg* 2018;**142**:734e–41e.
- Zeman-Kuhnert K, Gaggl AJ, Brandtner C, Wittig-Draenert AI, Bottini GB, Wittig J. Donor site morbidity after microvascular medial femoral condylar flap procurement for facial reconstruction. *Int J Oral Max Surg* 2020;**49**:569–75.
- Xia L, An J, He Y, Xiao E, Chen S, Yan Y, Zhang Y. Association between the clinical features of and types of temporomandibular joint ankylosis based on a modified classification system. *Sci Rep* 2019;**9**:10493.
- Binkley JM, Stratford PW, Lott SA, Riddle DL. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. North American Orthopaedic Rehabilitation Research Network. *Phys Ther* 1999;**79**:371–83.
- Mercuri LG. Avoiding and managing temporomandibular joint total joint replacement surgical site infections. *J Oral Maxillofac Surg* 2012;**70**:2280–9.
- Wolford LM, Rodrigues DB, McPhillips A. Management of the infected temporomandibular joint total joint prosthesis. *J Oral Maxillofac Surg* 2010;**68**:2810–23.
- Huang D, Lu C, Yao Z, He D, Yang C. A comparison of the effect between coronoid process graft and costochondral graft in the reconstruction of temporomandibular joint. *J Craniofac Surg* 2016;**27**:e197–200.
- Xia L, He Y, An J, Chen S, Zhang Y. Condyle-preserved arthroplasty versus costochondral grafting in paediatric temporomandibular joint ankylosis: a retrospective investigation. *Int J Oral Maxillofac Surg* 2019;**48**:526–33.
- Xia L, Zhang Y, An J, Chen S, He Y. Evaluating the remodeling of condyles reconstructed by transport distraction osteogenesis in the treatment of temporomandibular joint ankylosis. *J Craniomaxillofac Surg* 2020;**48**:494–500.
- Chen M, Yang C, Qiu Y, He D, Huang D, Wei W. Superior half of the sternocleidomastoid muscle for reconstruction of the temporomandibular joint: a preliminary study with a simplified technique and expanded indications. *Int J Oral Maxillofac Surg* 2015;**44**:685–91.
- Zaid WY, Alshehry S, Zakhary G, Yampolsky A, Kim B. Use of vascularized myo-osseous fibula free flap to reconstruct a hemimandibular defect with a concomitant skull defect arising from stock condylar prosthesis displacement into the middle cranial fossa. *J Oral Maxillofac Surg* 2019;**77**:1311–6.
- Lonie S, Herle P, Paddle A, Pradhan N, Birch T, Shayan R. Mandibular reconstruction: meta-analysis of iliac- versus fibula-free flaps. *Anz J Surg* 2016;**86**:337–42.
- Ling XF, Peng X. What is the price to pay for a free fibula flap? A systematic review of donor-site morbidity following free fibula flap surgery. *Plast Reconstr Surg* 2012;**129**:657–74.
- Gaggl AJ, Burger HK, Chiari FM. Free microvascular transfer of segmental corticocancellous femur for reconstruction of the alveolar ridge. *Br J Oral Maxillofac Surg* 2008;**46**:211–7.
- Dubois G, Lopez R, Puwanarajah P, Noyelles L, Lauwers F. The corticoperiosteal medial femoral supracondylar flap: anatomical study for clinical evaluation in mandibular osteoradionecrosis. *Surg Radiol Anat* 2010;**32**:971–7.
- Brandtner C, Hachleitner J, Bottini GB, Buerger H, Gaggl A. Microvascular medial femoral condylar flaps in 107 consecutive reconstructions in the head and neck. *Br J Oral Maxillofac Surg* 2016;**54**:614–8.
- Lee CC, Hackenberg B, Halvorson EG, Catterson EJ. Vascularized treatment options for reconstruction of the ascending mandible with introduction of the femoral medial epicondyle free flap. *J Craniofac Surg* 2014;**25**:1690–7.
- Higgins JP, Borumandi F, Burger HK, Benlidayi ME, Vasilyeva A, Sencar L, Polat S, Gaggl AJ. Nonvascularized cartilage grafts versus vascularized cartilage flaps: comparison of cartilage quality 6 months after transfer. *J Hand Surg Am* 2018;**43**:181–8.
- Benlidayi ME, Gaggl A, Buerger H, Kurkcu M, Unlugenc H, Onal D, Polat S, Sencar L. Comparison of vascularized osteoperiosteal femur flaps and nonvascularized femur grafts for reconstruction of mandibular defects: an experimental study. *J Oral Maxillofac Surg* 2009;**67**:1174–83.
- Kalickie T, Burger H, Muller EJ. A new vascularized cartilage-bone-graft for scaphoid nonunion with avascular necrosis of the proximal pole. Description of a new type of surgical procedure. *Unfallchirurg* 2008;**111**:201–5.
- Windhofer C, Wong VW, Larcher L, Paryavi E, Burger HK, Higgins JP. Knee donor site morbidity following harvest of medial femoral trochlea osteochondral flaps for carpal reconstruction. *J Hand Surg Am* 2016;**41**:610–4.
- Burger HK, Windhofer C, Gaggl AJ, Higgins JP. Vascularized medial femoral trochlea osteochondral flap reconstruction of advanced Kienbock disease. *J Hand Surg Am* 2014;**39**:1313–22.
- Higgins JP, Burger HK. Medial femoral trochlea osteochondral flap: applications for scaphoid and lunate reconstruction. *Clin Plast Surg* 2017;**44**:257–65.
- Iorio ML, Masden DL, Higgins JP. The limits of medial femoral condyle corticoperiosteal flaps. *J Hand Surg Am* 2011;**36**:1592–6.
- Hugon S, Koninckx A, Barbier O. Vascularized osteochondral graft from the medial femoral trochlea: anatomical study and clinical perspectives. *Surg Radiol Anat* 2010;**32**:817–25.
- Enzinger S, Bürger H, Gaggl A. Reconstruction of the mandibular condyle using the microvascular lateral femoral condyle flap. *Int J Oral Max Surg* 2018;**47**:603–7.

Address:

Yang He
Department of Oral and Maxillofacial Surgery
Peking University School and Hospital of Stomatology
22 Zhongguancun South Avenue
Haidian District
Beijing 100081
PR China
Tel.: +86 18911802149;
Fax: +86 10 82195158
E-mail: fridaydust1983@163.com