

STUDY PROTOCOL

Open Access



Assessment of soft and hard tissue characteristics of ridge preservation at molar extraction sites with severe periodontitis: a randomized controlled trial

Haoyun Zhang^{1†}, Yiping Wei^{1†}, Tao Xu², Min Zhen¹, Cui Wang¹, Ziyao Han¹, Wenjie Hu^{1*}  and Kwok-Hung Chung³

Abstract

Background: Changes in alveolar bone dimension after tooth extraction may affect placement of the subsequent implant, resulting in ridge deficiency that can adversely impact long-term implant stability or aesthetics. Alveolar ridge preservation (ARP) was effective in reducing the amount of ridge resorption following tooth extraction. There is sparse evidence regarding the benefit of ARP at periodontally compromised molar extraction sockets. This study will be a randomized trial to assess the soft tissue contour, radiographical, and histological changes of ARP at molar extraction sites in order to compare severe periodontitis cases with natural healing results and determine the most beneficial and least traumatic clinical treatment for such patients.

Methods: This research is designed as a two-group parallel randomized controlled trial. The total number of tooth extraction sites will be 70 after calculation with power analysis. Teeth will be randomly assigned to two groups with the test group conducting ridge preservation and the control group healing naturally. Periodontal examination, cone beam-computed tomography (CBCT) data, and stereolithographic (STL) files obtained by intraoral scanning will be collected through the follow-up period, and bone biopsy samples would be obtained during implant surgery. The primary outcomes are the vertical and horizontal change of alveolar ridge measured on CBCT images, soft tissue contour changes evaluated by superimposing the digital impressions, alterations of mucosa thickness (as measured by superimposing the CBCT data and STL files), histological features of implant sites and periodontal parameter changes. The secondary outcomes are patient-reported post-operative reaction and conditions of simultaneous bone graft or sinus lifting procedures during implantation.

[†]Haoyun Zhang and Yiping Wei contributed equally to this work.

*Correspondence: huwenjie@pkuss.bjmu.edu.cn

¹ Department of Periodontology, Peking University School and Hospital of Stomatology, National Center of Stomatology, National Clinical Research Center for Oral Diseases, National Engineering Laboratory for Digital and Material Technology of Stomatology, NHC Research Center of Engineering and Technology for Computerized Dentistry, 22 Zhongguancun S Ave, Haidian District, 100081 Beijing, China
Full list of author information is available at the end of the article



Discussion: This study will provide information about hard and soft tissue dimension changes and histomorphology evaluation following ARP and natural healing in periodontally compromised molar sites, which may contribute to complement the missing information of ARP at periodontally compromised molar extraction sockets.

Trial registration: Chinese Clinical Trial Register (ChiCTR) ChiCTR2200056335. Registered on February 4, 2022, Version 1.0.

Keywords: Severe periodontitis, Tooth extraction, Alveolar ridge preservation, Natural healing, Cone beam-computed tomography, Soft tissue contour, Histological evaluation

Background

Periodontitis is a chronic inflammatory condition which affects the tissues surrounding and supporting the teeth. Globally, severe periodontitis was the sixth-most prevalent health condition, affecting about 10.8% of population [1]. In recent decades, the total burden of periodontitis has increased [2], and it has become the main cause of tooth loss in adults [3]. Molar plays an important role in oral function, while it presents a challenge for periodontal treatment [4], and shows a higher risk for tooth loss due to its anatomy and position [5].

A systematic review reported that a certain amount of alveolar bone resorption occurs after natural healing, and more reduction can be expected in molar extraction sites [6]. This dimensional change of the alveolar ridge may affect the subsequent implant procedure, resulting in ridge deficiencies that can adversely impact long-term implant stability or aesthetics [7–9], necessitating additional reconstructive surgery [10, 11]. A consensus report concluded that alveolar ridge preservation (ARP) attenuated the bone resorption following tooth extraction [12]. Moreover, the indications for ARP procedures have been widening from anterior intact extraction socket to infected molar extraction sockets [13–16].


Regarding infected molar extraction, Zhao et al. [17] has confirmed that the molar extraction sockets with advanced periodontitis did resorb and change in various sites, especially at the buccal wall of the socket. Fok et al. [18] reported that prosthetically driven implants planning at first molar extraction sites due to terminal periodontitis poses greater challenge to rehabilitation, often requiring advanced augmentation procedures and sinus augmentation. Nevertheless, Kim et al. [19] revealed that ridge preservation at periodontally compromised sockets was safe and effective in reducing bone resorption. Wei et al. [20] concluded that ridge preservation could decrease the necessity of further regenerative procedures at maxillary molars with severe periodontitis during implant placement compared to natural healing sockets.

In measurement technique, most of the previous investigations usually used cone beam-computed tomography (CBCT) or peri-apical film to evaluate

the change of width and height of the alveolar socket received ARP [13, 20–23]. Only a few studies reported on the three-dimensional soft tissue profile changes using intraoral scanning technique for infected molar extraction sockets [24–26].

Histological analysis is valuable in assessing and determining the clinical application of ARP, since this procedure is often related to a following dental implant therapy that requires sufficient osseointegration within genuine bone tissue [27]. Previous studies showed ridge preservation using deproteinized bovine bone mineral (DBBM) had a lower amount of new bone formation in comparison with natural healing after a follow-up period of 3–6 months [28]. Further studies with longer healing time are needed for optimizing the clinical effects of ridge preservation in periodontally compromised sockets.

Although many previous studies have confirmed the effectiveness of ARP, few studies have been conducted in periodontally compromised molar extraction sites. Soft tissues surrounding implants have significant impact on esthetics and long-term health for dental implant therapy, but the comparison between natural healing and ARP in this area is also lacking. The different histological manifestations and the effect of implant therapy for these methods are also worthy of further exploration. Therefore, this randomized controlled trial is designed to compare the efficacy of ARP and natural healing at molar extraction sites in cases with severe periodontitis. This study is expected to determine the most beneficial and least traumatic clinical treatment for such patients. The following aspects will be compared, (i) soft tissue contour and mucosa thickness change from the baseline till at least 1 year after implant loading, (ii) radiographical evaluation for vertical and horizontal change of alveolar ridge from baseline to 6 months after atraumatic tooth extraction and ARP, (iii) histological composition after 6-month healing period in /tooth extraction sites with ARP, (iv) periodontal parameter changes in the follow-up period, (v) patient-reported post-operative reaction, and (vi) finally the need of simultaneous bone graft



or sinus lifting procedures during implant placement procedure.

Methods

This research is a two-group parallel randomized controlled trial (RCT) to provide soft tissue contour, radiographical, and histological evaluation of the efficacy of ARP in 70 molar extraction sites with severe periodontitis. Figure 1 shows the framework of this trial. The design and report of this protocol follow the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) statement [29], and the schedule of this study is shown in Fig. 2.

Setting

This trial will be undertaken at the Peking University School and Hospital of Stomatology (Beijing, China).

Before the trial takes place, written informed consent will be obtained from all participants.

Participants

If a patient has unsalvageable molars as a consequence of severe periodontal disease and considers implant-retained prostheses for the extraction sockets, the patient will be suggested and reclused in this study.

The included patient should meet the following criteria:

- 1) Age more than 25 years
- 2) Good compliance and good oral hygiene
- 3) Diagnosed with stage III/IV periodontitis [30]
- 4) Presence of at least one hopeless molar with severe bone loss requiring extraction
- 5) At least two socket walls beyond the apex or ≥ 3 mm of the extraction socket
- 6) At least one adjacent tooth at the proximal region.

TIMEPOINT	STUDY PERIOD														
	Enrolment	Allocation	Post-allocation												Close-out
	-t ₁	0	Surgery	1w	2 w	1 mo	3 mo	6 mo	Implant therapy	Implant loading	1 mo	3 mo	6 mo	Every 6 months after implant loading	Last appointment before December 31, 2024
ENROLMENT:															
Eligibility screen	x														
Informed consent	x														
Clinical examination	x														
SRP	x														
OHI	x														
Full-mouth periapical films	x														
Randomization and allocation		x													
INTERVENTIONS:															
ARP			x												
Atraumatical extraction			x												
Clinical maintenance								x		x			x	x	x
ASSESSMENTS:															
CBCT		x	x					x							
Intraoral scan		x	x		x	x	x	x	x	x	x	x	x	x	x
Bone biopsy									x						
WKT		x						x	x	x	x	x	x	x	x
PD		x						x	x	x	x	x	x	x	x
BI		x						x	x	x	x	x	x	x	x
GR		x						x	x	x	x	x	x	x	x
Post-operative reaction				x	x	x									
Simultaneous bone graft & sinus lifting procedures									x						

Fig. 2 Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) Figure: schedule of enrolment, interventions and assessments. SRP: scaling and root planning; OHI: Oral hygiene instruction; ARP: Alveolar ridge preservation; CBCT: Cone beam-computed tomography; WKT: Width of keratinized tissue; PD: Probing depth; BI: Bleeding index; GR: Gingival recession

The exclusion criteria are as followed:

- 1) Pregnancy or lactation
- 2) Patients taking medications or having disease that would complicate bone healing
- 3) Patients with surgical contraindications, such as uncontrolled hypertension or diabetes mellitus
- 4) Patients with history of head and neck radiotherapy
- 5) Smoking more than 10 cigarettes per day
- 6) Absence of both adjacent teeth
- 7) Teeth with ongoing acute pathology
- 8) Teeth extracted due to caries, endodontic failures or fractured teeth.

Criteria for discontinuing

The exclusion criteria will be checked regularly during the study period. If the participant meets any of the exclusion criteria during the follow-up period of this study, she/he will be excluded from the study. If the

participant shows poor compliance or decides to withdraw the consent due to any reason, the participation will be terminated.

Strategies to improve adherence

All time points in this study procedure will be matched with routine maintenance visits as far as possible to ensure the adherence to this clinical trial.

Randomizing

A series of random numbers will be generated using Excel 2021 (Microsoft Corp., Redmond, WA, USA), written on letter paper and sealed in sequentially numbered, sealed, opaque envelopes. All the patients will be assigned to two groups by simple randomization with the rate of 1:1: the test group with alveolar ridge preservation, and the control group with natural healing. All the surgical procedures will be performed by an experienced clinician, and the other clinician will conduct the clinical examination for the patients. Another two members will take charge of intraoral scanning, assessing

the radiological examinations and completing statistical analysis.

Blinding

Blinding is not suitable for this trial. For the participants, they can distinguish which group they are in after experiencing different surgical procedures and post-operative treatment, so can the surgeons. For the assessors, they can also differentiate two groups from different imaging manifestations.

Intervention

The enrolled patients will receive clinical and radiological examinations. They will receive routine scaling and root planing, and oral hygiene instruction at least 1 week before the surgery. All the patients will also receive prophylactic antibiotic medicine (Amoxicillin, 1 g or Erythromycin 300 mg if allergic to Penicillin) and anti-inflammatory drug (Ibuprofen 300 mg) 1 h prior to the surgical procedures described previously [31].

For control group:

- (1) An internal bevel incision will be performed from 0.5–1.0 mm below the buccal and lingual gingival margin to the bone crest.
- (2) The selected tooth will be extracted atraumatically [31].
- (3) The socket will be examined carefully and debrided thoroughly (using P24G Periosteal Elevator and CL86 Lucas Surgical Curette) without damaging the socket walls.

For test group:

- (1) The steps mentioned above are the same.
- (2) A full thickness flap will be elevated to expose 2 mm of the bone crest.
- (3) The socket will be filled with DBBM (Bio-Oss, Geistlich Pharma AG, Wolhusen, Switzerland), and a resorbable membrane (Bio-Gide, Geistlich Pharma AG, Wolhusen, Switzerland) will be used to cover the socket completely with 2 mm extending over the crest.
- (4) Then the socket will be covered by collagen sponge (Wuxi Biot Biology Technology Co.,Ltd., Wuxi, China).
- (5) A cross-mattress tension-free 5–0 suture will be placed over the site.

After the surgical procedures, patients will be prescribed to take the antibiotic three times daily for 7 days and ibuprofen (300 mg twice daily for 3–5 days)

if needed. Patients will be required to perform regular tooth brushing in the rest of the mouth and oral rinse with 0.12% chlorhexidine solution for the 4-week post-surgery. The sutures will be removed 2 weeks after the surgery, and all the patients will be recalled at 4 weeks after the surgical to evaluate the healing status of the surgery site.

During the whole follow-up period from baseline to the last appointment, all the participants will receive regular maintenance therapy.

Outcome measures


Baseline examination (Fig. 3)

At baseline (before tooth extraction), the following data will be collected:

- (1) Full-mouth peri-apical films,
- (2) CBCT images (CBCT 1): CBCT images will be taken by CBCT machines (NewTom VG; Aperio Services, Italy) at a resolution of 0.125 mm with field of view size 10 × 10 cm (exposure time: 3.6 s, 110 kV, 5 mA). It will be used for assessing anatomical conditions, including the degree of bone loss and the position of maxillary sinus/mandibular canal.
- (3) Stereolithography (STL) files (STL 1): intraoral scans will be taken using a TRIOS intraoral scanner (3Shape TRIOS Color, TRC, 3Shape, Denmark) to obtain a digital impression of the target area and generate an STL file, which include the selected tooth and relevant quadrant. Soft tissues contained in this digital impression will extend over the mucogingival junction.
- (4) Periodontal parameters: the periodontal conditions of the ailing tooth and adjacent teeth will be assessed by measuring probing depth (PD), gingival recession (GR) and bleeding index (BI). The width of keratinized tissue (WKT) will be measured at mid-buccal aspect from the mucogingival junction to the gingival margin of the teeth. Data will be round down to the nearest 0.5 mm, and they will be collected using an UNC-15 probe (Hu-Friedy, Chicago, IL, USA).

Examination immediately after the intervention for both groups (Fig. 3)

- (1) CBCT images (CBCT 2): it will be used for superimposition and measurement. And it will be illustrated in informed consent.
- (2) STL files (STL 2)



Examinations during the follow-ups before implant therapy (Fig. 3)

- (1) STL files (STL 3–6): the patients will be recalled at 2, 4, 12 weeks and 6 months after the surgery, and intraoral scans will be performed.
- (2) CBCT images (CBCT 3): CBCT examination will be performed 6 months after the surgery.
- (3) Post-operative reaction: the patients will be recalled at 1, 2 and 4 weeks after the surgery to report post-operative reaction, such as pain, erythema, swelling and paranesthesia.
- (4) Paralleling technique films: films will be taken 6 months after the surgery.
- (5) Periodontal parameters: at 6 months after the surgery, the periodontal examination indicators (PD, GR, BI and WKT) will be recorded for the adjacent teeth, and WKT of extraction area will also be measured at the buccal aspect from the central of the (expected) implant position to the buccal mucogingival junction [32].

Examination during implant therapy (Fig. 3)

The implant therapy will be performed around 6 months after ARP/tooth extraction.

- 1) Bone biopsy: during the implant placement procedure, a cylindrical bone block with a diameter of about 2 mm and length of about 5 mm will be taken out using a trephine drill (Institute Straumann AG, Basel, Switzerland). The block will be fixated in 4% formaldehyde solution at room temperature for at least one day. Then, they will receive decalcification for 3 weeks in 4.1% disodium ethylenediamine-tetra-acetic acid-solution, the solution will be changed every 24 h. After hydration, they will be dehydrated in increasing concentrations of ethanol (from 70 to 100%), embedded in paraffin, and cut into sections with a thickness of 4 μ m. Serial sagittal sections will be stained with hematoxylin–eosin. Representative regions of interest (ROIs) will be localized in the center, apically and coronally of the sample [33]. Histologic slides will be observed under light microscopes and digital images will be obtained by scanning at a magnification of $\times 200$ for subsequent histomorphometric analysis. For the two groups, the histological composition in terms of new bone formation, residual graft particles and connective tissue will be evaluated and compared.
- 2) Simultaneous bone graft & sinus lifting procedures: the conditions of bone grafting and sinus lifting during implant therapy will be recorded.

Examination during the follow-ups after implant restoration (Fig. 3)

After 6-month period of osseointegration, the superstructures will be connected to the implants. Participants will be recalled at immediately, 1-, 3-, and 6-month and then every 6 months after implant loading. During this period, the following data will be collected:

- 1) STL files (STL R1, R2, and so on): intraoral scans will be performed at every appointment.
- 2) Periodontal parameters: the periodontal examination indicators (PD, GR, BI and WKT) will be recorded for the implant and adjacent teeth at every appointment.
- 3) Paralleling technique films: the films will be collected immediately and every 6 months after implant loading, and they will be used to evaluate the marginal bone loss (MBL).

Superimposition and measurement of CBCT images and digital impressions

- (1) CBCT images: Digital Imaging and Communications in Medicine (DICOM) data will be generated and transferred to a volumetric imaging software (Mimics 21.0, Materialise, Leuven, Belgium). Residual buccal and lingual (or palatal) plate thicknesses and height of residual socket walls can be measured and recorded with vertical/horizontal reference lines and dots on CBCT 2. After Superimposition of CBCT 2 and CBCT 3, horizontal width and vertical height changes at target sites will be calculated; details are discussed in the previous literature [13, 17].
- (2) STL files: these data will be generated and imported to Geomagic Studio 2021 (3D Systems Inc., Rock Hill, SC, USA). Superimposition of STL 1 and STL files for subsequent stages (STL 2–6, STL R1, and so on) will be completed for the selected areas, using identified landmarks such as cusps of adjacent teeth. After superimposition, the impressions can be aligned and manually checked for perfect matching, and changes of the contour lines of alveolar ridge can be observed and measured.
- (3) CBCT data and STL files: As described in previous literature [25], CBCT 2 and STL 3 will be superimposed in Geomagic Studio 2021 (3D Systems Inc., Rock Hill, SC, USA) using three or more landmarks of adjacent teeth and so will CBCT 3 and STL 6. The vertical thickness of the crestal mucosa and

horizontal thickness of the buccal and palatal/lingual mucosa and their alterations will be measured.

Sample size calculation

The sample size was calculated using the Power Analysis and Sample Size (PASS) (version 15.0, NCSS, LLC, East Kaysville, Utah, USA) based on a two-sided alpha of 0.05 and power of 0.8. According to previous literature [13], the ridge width changes in one site (at 1 mm apically from the top of the alveolar crest in central buccal) of ridge preservation group and natural healing group were 1.46 mm and -0.70 mm, respectively. And the SD were 3.54 mm and 2.28 mm. After the calculation, the minimal sample size was 30 extraction sites for each group. Considering 15% non-response rate, 35 extraction sites will be included in this study for each group. Totally, 70 extraction sites will be needed.

Statistical analysis

Statistical analysis will be performed using the SPSS 26.0 software package (SPSS Inc., Chicago, IL, USA). Descriptive data will be reported as mean \pm standard deviations. A Shapiro–Wilk test will be applied to test for normal distribution of the sample for each variable. The paired t-test or Wilcoxon signed rank sum test when the data are not normally distributed will be applied to detect changes of soft and hard tissue before and 6 months after surgery. Independent samples t tests will be performed to compare parameter means between control and test groups. A non-parametric Mann–Whitney U test will be used if parameters are not normally distributed. Baseline characteristics of the patients who are lost to follow-up and completed this study will be compared. For the missing outcome data, K-means clustering method will be used for imputation. The level of significance was set at $\alpha = 0.05$.

Research ethics committee/institutional review board (REC/IRB) approval

This trial will be performed in strict accordance with the World Medical Association Declaration of Helsinki. The study protocol and written documents has been reviewed by the Ethics Committee of Peking University School and Hospital of Stomatology and received approval (PKUS-SIRB-202170189). Written informed consent to participate will be obtained from all participants.

Withdrawal

Participants will be informed that they can quit this study at any time without influencing their treatment process in the future.

Discussion

This is a two-group parallel randomised controlled trial to compare the multi-dimensional efficacy of ARP and natural healing at molar extraction sites with severe periodontitis. Since Kim et al. [19] showed the safety and effectiveness of ridge preservation at periodontally compromised sockets, an increasing number of studies has reported that ridge preservation in these sites was effective in reducing the amount of ridge resorption [13, 14, 20]. Wei et al. [20] evaluated ridge preservation in the maxillary molar extraction sockets with severe periodontitis, and concluded that ARP could decrease the necessity of advanced regenerative procedures at implant placement compared to natural healing. A recent retrospective study included 418 extraction sites (171 without ARP and 247 with ARP) also showed ARP may reduce the need for bone augmentation procedure and improve the feasibility of implant placement [16]. But most of these studies merely used dental records and imaging data to evaluate the efficacy.

According to Avila-Ortiz et al. [34], peri-implant phenotype is crucial to peri-implant health, function, and esthetics. Soft tissues take over an important position in the components of peri-implant phenotype. The use of intraoral scan allows to record the profile of soft tissues without any pressure. It can be used immediately after surgical procedures without disrupting the wound. Digital scanned data make it possible to measure the dimensional changes at different time points in a long-term follow-up period [26]. Up to now, only a few studies reported the three-dimensional profilometric changes on the soft tissue level based on intraoral scanned data following tooth extraction and ARP [24–26]. In the esthetic zone of single extraction sites, Chappuis et al. [24] investigated the interplay between the soft tissue morphology and the underlying bone anatomy during an 8-week healing period by sequential digitized impressions and CBCT images. Wongpairojpanich et al. [26] used intraoral scanner for monitoring the wound closure and surface dimensional changes of alveolar ridge at different time points (1, 3, 7, 14, and 28 days and 4 months after ARP). Song et al. [25] explored the changes of soft tissue dimensions following alveolar ridge preservation and spontaneous healing in posterior maxilla using intraoral scanned data and CBCT data in a 6-month healing period. To summarize, the follow-up time for current studies was relatively short. And to the best of our knowledge, there has not been specific evidence available to identify the soft tissue contour changes of ridge preservation on molar extraction sockets with periodontal pathosis before tooth extraction.

DBBM is a type of bone substitute material widely used in dental practice [35]. A systematic review included five studies showed that ridge preservation using DBBM had a lower amount of new bone formation in comparison with natural healing, but due to the lack of randomized controlled trials, limited sample size and high heterogeneity, further experiments are needed to verify this conclusion [28]. Koo et al. [36] carried out a histological analysis at 4-month after ARP in damaged extraction sites, showing a low mean proportion of regenerated bone and a large variation from 0 to 43%. Further studies with longer healing time in periodontally compromised sockets are needed for evaluating the histological changes of ridge preservation.

This study will produce data on hard and soft tissue dimension changes at different time points and provide histomorphology evaluation following ARP and natural healing in periodontally compromised molar extraction sockets. The follow-up period of this trial will last till at least 1 year after implant loading. These results may contribute to complement the missing information for ARP in molar sites with advanced periodontitis. Results of the study will be published in peer-reviewed journals.

Trial registration

The trial has been registered at Chinese Clinical Trial Register (ChiCTR) with the identifier number ChiCTR2200056335 on February 4, 2022. The recruitment began in March 1, 2022 and will be completed in March 1, 2023.

Abbreviations

ARP: Alveolar ridge preservation; BI: Bleeding index; CBCT: Cone beam-computed tomography; DBBM: Deproteinized bovine bone mineral; DICOM: Digital Imaging and Communications in Medicine; GR: Gingival recession; MBL: Marginal bone loss; OHI: Oral hygiene instruction; PD: Probing depth; RCT: Randomised controlled trial; ROIs: Representative regions of interest; SRP: Supra-/sub-gingival scaling, root planning; STL: Stereolithographic; WKT: Width of keratinized tissue.

Acknowledgements

The authors thank Dr. Joelle Moussi for her critical evaluation of this manuscript.

Authors' contributions

YW, HZ, TX and WH proposed the concept of this work. HZ, YW and WH designed the trial. HZ and YW drafted the document. MZ, CW and ZH revised the part on intervention and examination. WH and KC reviewed and finalised the manuscript. All authors agreed with the final version.

Funding

This work was supported by Capital's Funds for Health Improvement and Research, Grant # CFH 2022–2-4103. The funders had no role in study design, data collection and analysis, or manuscript preparation.

Availability of data and materials

Data from this research will be registered with the Chinese Clinical Trial Registry Platform. Results will be published through official peer-reviewed journals.

Declarations

Ethics approval and consent to participate

The present research has received approval from the Ethics Committee of Peking University School and Hospital of Stomatology (PKUSSIRB-202170189). Written informed consent to participate will be obtained from all participants.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Periodontology, Peking University School and Hospital of Stomatology, National Center of Stomatology, National Clinical Research Center for Oral Diseases, National Engineering Laboratory for Digital and Material Technology of Stomatology, NHC Research Center of Engineering and Technology for Computerized Dentistry, 22 Zhongguancun S Ave, Haidian District, 100081 Beijing, China. ²Department of Emergency, Peking University School and Hospital of Stomatology, National Center of Stomatology, National Clinical Research Center for Oral Diseases, National Engineering Laboratory for Digital and Material Technology of Stomatology, NHC Research Center of Engineering and Technology for Computerized Dentistry, Beijing, China. ³Department of Restorative Dentistry, School of Dentistry, University of Washington, Seattle, WA, USA.

Received: 5 August 2022 Accepted: 27 October 2022

Published online: 17 November 2022

References

- Peres MA, Macpherson LMD, Weyant RJ, Daly B, Venturelli R, Mathur MR, et al. Oral diseases: a global public health challenge. *Lancet*. 2019;394(10194):249–60. [https://doi.org/10.1016/S0140-6736\(19\)31146-8](https://doi.org/10.1016/S0140-6736(19)31146-8).
- Jepsen S, Blanco J, Buchalla W, Carvalho JC, Dietrich T, Dorfer C, et al. Prevention and control of dental caries and periodontal diseases at individual and population level: consensus report of group 3 of joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J Clin Periodontol*. 2017;44:S85–93. <https://doi.org/10.1111/jcpe.12687>.
- Frencken JE, Sharma P, Stenhouse L, Green D, Laverty D, Dietrich T. Global epidemiology of dental caries and severe periodontitis - a comprehensive review. *J Clin Periodontol*. 2017;44(Suppl 18):S94–105. <https://doi.org/10.1111/jcpe.12677>.
- Dannewitz B, Zeidler A, Husing J, Saure D, Pfefferle T, Eickholz P, et al. Loss of molars in periodontally treated patients: results 10 years and more after active periodontal therapy. *J Clin Periodontol*. 2016;43(1):53–62. <https://doi.org/10.1111/jcpe.12488>.
- Baumer A, Pretzl B, Cosgarea R, Kim TS, Reitmeir P, Eickholz P, et al. Tooth loss in aggressive periodontitis after active periodontal therapy: patient-related and tooth-related prognostic factors. *J Clin Periodontol*. 2011;38(7):644–51. <https://doi.org/10.1111/j.1600-051X.2011.01733.x>.
- Couso-Queiruga E, Stuhr S, Tattan M, Chambrone L, Avila-Ortiz G. Post-extraction dimensional changes: A systematic review and meta-analysis. *J Clin Periodontol*. 2021;48(1):126–44. <https://doi.org/10.1111/jcpe.13390>.
- Buser D, Martin W, Belsler UC. Optimizing esthetics for implant restorations in the anterior maxilla: anatomic and surgical considerations. *Int J Oral Maxillofac Implants*. 2004;19(Suppl):43–61.
- Van der Weijden F, Dell'Acqua F, Slot DE. Alveolar bone dimensional changes of post-extraction sockets in humans: a systematic review. *J Clin Periodontol*. 2009;36(12):1048–58. <https://doi.org/10.1111/j.1600-051X.2009.01482.x>.
- Tan WL, Wong TLT, Wong MCM, Lang NP. A systematic review of post-extraction alveolar hard and soft tissue dimensional changes in humans. *Clin Oral Implan Res*. 2012;23:1–21. <https://doi.org/10.1111/j.1600-0501.2011.02375.x>.
- Hu C, Gong T, Lin WM, Yuan Q, Man Y. Immediate implant placement into posterior sockets with or without buccal bone dehiscence defects: A retrospective cohort study. *J Dent*. 2017;65:95–100. <https://doi.org/10.1016/j.jdent.2017.07.010>.
- Milinkovic I, Cordaro L. Are there specific indications for the different alveolar bone augmentation procedures for implant placement? A systematic review. *Int J Oral Max Surg*. 2014;43(5):606–25. <https://doi.org/10.1016/j.ijom.2013.12.004>.
- Tonetti MS, Jung RE, Avila-Ortiz G, Blanco J, Cosyn J, Fickl S, et al. Management of the extraction socket and timing of implant placement: Consensus report and clinical recommendations of group 3 of the XV European Workshop in Periodontology. *J Clin Periodontol*. 2019;46(Suppl 21):183–94. <https://doi.org/10.1111/jcpe.13131>.
- Zhao LP, Xu T, Hu WJ, Chung KH. Preservation and augmentation of molar extraction sites affected by severe bone defect due to advanced periodontitis: a prospective clinical trial. *Clin Implant Dent R*. 2018;20(3):333–44. <https://doi.org/10.1111/cid.12585>.
- Ben Amara H, Kim JJ, Kim HY, Lee J, Song HY, Koo KT. Is ridge preservation effective in the extraction sockets of periodontally compromised teeth? A randomized controlled trial. *J Clin Periodontol*. 2021;48(3):464–77. <https://doi.org/10.1111/jcpe.13412>.
- Kim JJ, Ben Amara H, Chung I, Koo KT. Compromised extraction sockets: a new classification and prevalence involving both soft and hard tissue loss. *J Periodontal Implant Sci*. 2021;51(2):100–13. <https://doi.org/10.5051/jpis.2005120256>.
- Lee J, Yun J, Kim JJ, Koo KT, Seol YJ, Lee YM. Retrospective study of alveolar ridge preservation compared with no alveolar ridge preservation in periodontally compromised extraction sockets. *Int J Implant Dent* 2021, 7(1) <https://doi.org/10.1186/s40729-021-00305-2>.
- Zhao L, Wei Y, Xu T, Zhang B, Hu W, Chung KH. Changes in alveolar process dimensions following extraction of molars with advanced periodontal disease: A clinical pilot study. *Clin Oral Implants Res*. 2019;30(4):324–35. <https://doi.org/10.1111/clr.13418>.
- Fok MR, Pelekos G, Tonetti MS. Feasibility and needs for simultaneous or staged bone augmentation to place prosthetically guided dental implants after extraction or exfoliation of first molars due to severe periodontitis. *J Clin Periodontol*. 2020;47(10):1237–47. <https://doi.org/10.1111/jcpe.13344>.
- Kim JJ, Ben Amara H, Schwarz F, Kim HY, Lee JW, Wikesjo UME, et al. Is ridge preservation/augmentation at periodontally compromised extraction sockets safe? A retrospective study. *J Clin Periodontol*. 2017;44(10):1051–8. <https://doi.org/10.1111/jcpe.12764>.
- Wei Y, Xu T, Zhao L, Hu W, Chung K-H. Ridge preservation in maxillary molar extraction sites with severe periodontitis: a prospective observational clinical trial. *Clin Oral Invest*. 2022;26(3):2391–9. <https://doi.org/10.1007/s00784-021-04204-z>.
- Lombardi T, Bernardello F, Berton F, Porrelli D, Rapani A, Camurri Piloni A, et al. Efficacy of Alveolar Ridge Preservation after Maxillary Molar Extraction in Reducing Crestal Bone Resorption and Sinus Pneumatization: A Multicenter Prospective Case-Control Study. *Biomed Res Int*. 2018;2018:9352130. <https://doi.org/10.1155/2018/9352130>.
- Zhao LP, Zhan YL, Hu WJ, Xu T, Wei YP, Zhen M, et al. Evaluation with different measuring methods for the alveolar bone change of ridge preservation in molar sites. *Beijing Da Xue Xue Bao Yi Xue Ban*. 2016;48(1):126–32. <https://doi.org/10.3969/j.issn.1671-167X.2016.01.023>.
- Walker CJ, Prihoda TJ, Mealey BL, Lasho DJ, Noujeim M, Huynh-Ba G. Evaluation of Healing at Molar Extraction Sites With and Without Ridge Preservation: A Randomized Controlled Clinical Trial. *J Periodontol*. 2017;88(3):241–9. <https://doi.org/10.1902/jop.2016.160445>.
- Chappuis V, Engel O, Shahim K, Reyes M, Katsaros C, Buser D. Soft Tissue Alterations in Esthetic Postextraction Sites: A 3-Dimensional Analysis. *J Dent Res*. 2015;94(9 Suppl):1875–1935. <https://doi.org/10.1177/0022034515592869>.
- Song YW, Yoon SW, Cha JK, Jung UW, Jung RE, Thoma DS. Soft Tissue Dimensions Following Tooth Extraction in the Posterior Maxilla: A Randomized Clinical Trial Comparing Alveolar Ridge Preservation to Spontaneous Healing. *J Clin Med* 2020, 9(8). <https://doi.org/10.3390/jcm9082583>.
- Wongpairorpanich J, Kijartorn P, Suwanprateeb J, Buranawat B. Effectiveness of bilayer porous polyethylene membrane for alveolar ridge preservation: a randomized controlled trial. *Clin Implant Dent R*. 2021;23(1):73–85. <https://doi.org/10.1111/cid.12965>.
- Guglielmotti MB, Olmedo DG, Cabrini RL. Research on implants and osseointegration. *Periodontol* 2000 2019, 79(1):178–189. <https://doi.org/10.1111/prd.12254>.

28. Zhao H, Hu J, Zhao L: Histological analysis of socket preservation using DBBM. A systematic review and meta-analysis. *J Stomatol Oral Maxi* 2020, 121(6):729–735. <https://doi.org/10.1016/j.jormas.2020.04.011>.
29. Chan AW, Tetzlaff JM, Gotzsche PC, Altman DG, Mann H, Berlin JA, et al. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *BMJ*. 2013;346:e7586. <https://doi.org/10.1136/bmj.e7586>.
30. Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, et al. Periodontitis: Consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Periodontol*. 2018;89(Suppl 1):S173–82. <https://doi.org/10.1002/JPER.17-0721>.
31. Wei Y, Xu T, Hu W, Zhao L, Wang C, Chung KH: Socket Preservation Following Extraction of Molars with Severe Periodontitis. *Int J Periodontics Restorative Dent* 2021, 41(2):269–275. <https://doi.org/10.11607/prd.4444>.
32. Han ZY, Wei YP, Wang C, Yang G, Hu WJ, Chung KH. Clinical evaluations of free gingival grafting before implant placement to increase keratinized tissue width in molar regions: A retrospective case series. *Clin Oral Implan Res*. 2021;32(7):799–807. <https://doi.org/10.1111/clr.13748>.
33. Schnutenhaus S, Edelmann C, Dreyhaupt J, Rudolph H, Luthardt RG, Goetz W. Alveolar ridge preservation with a collagen cone: Histological, histochemical, and immunohistochemical results of a randomized controlled clinical trial. *Clin Exp Dent Res*. 2020;6(3):345–55. <https://doi.org/10.1002/cre2.279>.
34. Avila-Ortiz G, Gonzalez-Martin O, Couso-Queiruga E, Wang H-L. The peri-implant phenotype. *J Periodontol*. 2020;91(3):283–8. <https://doi.org/10.1002/JPER.19-0566>.
35. Wessing B, Lettner S, Zechner W: Guided Bone Regeneration with Collagen Membranes and Particulate Graft Materials: A Systematic Review and Meta-Analysis. *Int J Oral Max Impl* 2018, 33(1):87–100. <https://doi.org/10.11607/jomi.5461>.
36. Koo TH, Song YW, Cha JK, Jung UW, Kim CS, Lee JS. Histologic analysis following grafting of damaged extraction sockets using deproteinized bovine or porcine bone mineral: A randomized clinical trial. *Clin Oral Implan Res*. 2020;31(1):93–102. <https://doi.org/10.1111/clr.13557>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

